

Dupuytren's Fasciectomy (for Dupuytren's Contracture) Surgery Guide

General Information

- You can eat and drink when you feel able to.
- Swelling and pain is normal, especially for the first 48 hours. Each patient experiences discomfort to a different degree but you can expect your discomfort to decrease rapidly over the first 48-72 hours postoperatively.
- Ice can be applied as needed for 20 minutes at a time, with 20 minute breaks in-between.
- Heat may be used as needed after the swelling is completely gone.
- Keep your arm elevated when not in use to reduce swelling. You may lower your arm for use, but this could cause your arm to throb and swelling to increase.
- Your doctor will give you a prescription for pain medication before you leave the hospital. Please check with us if you are allergic to any pain medication.
- You will have receive a local anesthetic during your operative procedure. In general, the local anesthesia continues to work for 4-12 hours postoperatively. This local anesthesia will greatly diminish your discomfort. However, as the anesthetic begins to wear off, you are encouraged to take your pain medication to help alleviate the ensuing anticipated discomfort.
- If you are experiencing any of the following, call our office or the doctor on call immediately:
 - o Fever greater than 100.4 °F
 - o Chills
 - o Shortness of breath
 - o Excessive redness or swelling of incisions
 - o Yellow (pus-like) or white drainage at incision sites
 - o Bleeding that soaks your dressings and cannot be controlled by firm pressure
 - o Pain not relieved by prescribed medications

How to reach your provider:

MyChart:

This is the easiest and quickest way to contact your provider. You may send a message anytime, and can include photos as needed.

By phone:

Weekdays 8 a.m. - 4:30 p.m.
Call the Upper Extremity
Orthopaedic Clinic at
503-494-8645.

After business hours/weekend/
holidays: Call the OHSU Hospital
Operator at 503-494-8311 and
ask to be connected to the on-call
orthopaedic resident.

www.ohsu.edu/ortho





Activity

- Limit weight-bearing activities to two pounds on the operative side. There are no restrictions with your range of motion.
- When you can return to work is dependent upon the type of work you do. Please ask your surgeon for specific work restrictions.
- You can drive once you are fully weight-bearing and no longer taking narcotics.

Wound Care

- It is OK to shower immediately following your procedure but please keep your wound clean and dry.
- No soaking or swimming until your wound has fully closed.

Occupational Therapy

- Occupational therapy to start approximately one to three days post-surgery after your first post-operative appointment with our clinic.
- Occupational therapy can take place at OHSU, or with an Occupational Therapist of your choice. Please let our office know where you are planning to go if outside of OHSU before your procedure.

Follow-up appointments you should have scheduled:

- You will either be scheduled for an occupational therapy appointment at OHSU one to three days after your procedure, or you will need to set-up an appointment with an outside Occupational Therapist to start one to three days after your procedure.

Scheduled on:

__/__/__ at __:__am/pm

- You will have your first post-operative appointment with our office approximately two weeks after your procedure with one of your surgeon's Physician Assistants. Your sutures will be removed at this appointment.

Scheduled on:

__/__/__ at __:__am/pm

- Your second post-operative appointment will be approximately six weeks after your procedure and will be with your surgeon. You will get x-rays at this appointment.

Scheduled on:

__/__/__ at __:__am/pm

Please let our office know if you need to make any of these appointments.